VOL. 520 JOURNAL OF CHROMATOGRAPHY NOVEMBER 9, 1990

CONTENTS

7TH DANUBE SYMPOSIUM ON CHROMATOGRAPHY, LEIPZIG, AUGUST 21–25, 1989

Foreword
by W. Engewald 1

GAS CHROMATOGRAPHY
Columns and sorbents

Enhanced possibilities for identification by the use of series-coupled capillary gas chromatographic columns. I. General exposition and application of the retention index concept
by W. Engewald and T. Maurer (Leipzig, G.D.R.) 3

Lead selenide as a polar adsorbent for gas chromatography
by T. M. Roshchina and E. V. Vasil'eva (Moscow, U.S.S.R.) 15

Polymer–inorganic selective adsorbents for gas chromatography produced by graft polymerization
by A. V. Dmitrenko, N. E. Shadrina, S. S. Ivanchev, N. N. Ulinskaya and A. M. Volkov (Leningrad, U.S.S.R.) 21

Short columns for gas chromatography packed with plasma-modified silica gels
by Chr. Chanev, N. Petsev, I. Topalova and St. Ivanov (Sofia, Bulgaria) and T. B. Gavrilova (Moscow, U.S.S.R.) 33

Gas chromatographic properties of modified boron nitride
by T. B. Gavrilova and I. V. Daidakova (Moscow, U.S.S.R.) and N. Petsev, I. Topalova, St. Ivanov and Chr. Chanev (Sofia, Bulgaria) 41

Retention and heats of adsorption of *n*-alkanes, aromatic hydrocarbons, aliphatic alcohols and ketones on macroporous silicas
by L. I. Dernovaya and Yu. A. Eltekov (Moscow, U.S.S.R.) 47

Carbon adsorbents in gas adsorption chromatography
by N. V. Kovaleva and K. D. Shcherbakova (Moscow, U.S.S.R.) 55

Determination of heats of adsorption on graphite and on a microporous carbon black by gas–solid chromatography
by S. Ludwig and H.-D. Schmidt (Berlin, G.D.R.) 69

High-resolution gas chromatography with liquid crystal glass capillaries. XII. Separation of isomeric C_{17}–C_{18} *n*-alkenes
by L. Soják, I. Ostrovský and R. Kubinec (Bratislava, Czechoslovakia) and G. Kraus and A. Kraus (Halle/Saale, G.D.R.) 75

Blended liquid crystalline stationary phases in capillary gas chromatography
by F. Hahne, G. Kraus and H. Zaschke (Halle/Saale, G.D.R.) 85

General

Chromatographic characteristics of capillary packed columns with increased pressure drop
by V. G. Berezkin, V. S. Gavrichev and N. V. Voloshina (Moscow, U.S.S.R.) 91

Radiofrequency detector for gas chromatography
by P. Popp, S. Mothes and D. Hinneburg (Leipzig, G.D.R.) 97

Chromatographic study of the interactions of hydrocarbons with cyclodextrins and their derivatives in the gas–solid system
by L. Feltl, P. Tajovská and E. Smolková-Keulemansová (Prague, Czechoslovakia) . . . 105

Chromatograms of second-order reactions on chromatographic columns
by R. Thede, E. Below, D. Haberland and H. Pscheidl (Greifswald, G.D.R.) 109

Probability method for pyrolysis gas chromatographic data treatment and elaboration of a computer system for the identification of polymers
by V. M. Ryabikova, S. D. Tumina and A. N. Zigel (Leningrad, U.S.S.R.) 121

Thermal desorption of organic pollutants enriched on activated carbon. V. Desorption behaviour and temperature profile
by L. Senf and H. Frank (Gera, G.D.R.) 131

Relationship between the structure and retention of *n*-alkenes and *n*-alkynes on silicone phases
by K. Kuningas, S. Rang and T. Kailas (Tallinn, U.S.S.R.) 137

Applications

Gas chromatographic–mass spectrometric identification of impurities in 1,2-diethoxybenzene
by I. Hazai and L. Ledniczky (Budapest, Hungary) 149

Investigation of phase transitions in the $CaSO_4(s)$–$H_2O(g)$ system by chromatographic methods
by L. G. Berezkina, V. I. Souhodolova, S. V. Melnikova and E. A. Saricheva (Moscow, U.S.S.R.) 157

Determination of impurities in thionyl chloride by gas chromatography
by W. Czerwinski and W. Gromotowicz (Warsaw, Poland) 163

Headspace gas chromatography with open-tubular columns for determining the evaporation rate of essential oils, perfumery compositions and perfumes
by M. A. Dubrova, L. O. Mindlin, M. M. Shchedrina, A. G. Belfer, M. A. Nikitina and B. A. Rudenko (Moscow, U.S.S.R.) 169

Packed column gas chromatography–capillary column gas chromatography–mass spectrometry coupling for the qualitative characterization of technical products
by D. Estel, J. Ohl and C. E. Döring (Leuna, G.D.R.) 175

Gas–liquid chromatographic assay of the components of electrochemically reduced glucose solutions. Impurities in sorbitol samples
by I. Molnár-Perl, M. Morvai and M. Pintér-Szakács (Budapest, Hungary) and Á. Kővágó and J. Petróczy (Nádudvar, Hungary) 185

Gas chromatography of tryptophan together with other amino acids in hydrochloric acid hydrolysates
by V. Fábián, M. Pintér-Szakács and I. Molnár-Perl (Budapest, Hungary) 193

Simultaneous determination of organic acids and sugars in apples by gas–liquid chromatography
by M. Morvai and I. Molnár-Perl (Budapest, Hungary) 201

Speciation of polyhydroxy compounds by gas chromatography–mass spectrometry with deuterium oxide chemical ionization (Note)
by M. Svoboda and S. Musil (Modra, Czechoslovakia) 209

Analysis of internal filling gas of small-sized halogen lamps by gas chromatography
by L. Nagy, E. Biro and J. Balla (Budapest, Hungary) 213

Reversed-phase thin-layer and gas chromatographic retention behaviour of triphenylmethane derivatives
by E. János and G. Oros (Budapest, Hungary) 217

LIQUID CHROMATOGRAPHY

General

Feather keratin as a ligand in an affinity chromatographic technique for isolation of protease from *Trichophyton verrucosum*
by J. Łobarzewski, K. Grzywnowicz, K. Wawrzkiewicz, M. Staszczak and T. Wolski (Lublin, Poland) 223

Studies on texture and properties of porous glass and silica gel used as supports for chemically bonded amino phases for high-performance liquid chromatography
by B. Buszewski, M. Hummel and E. Bayer (Tübingen, F.R.G.) 237

Determination of the lipophilicity of some peptides. Effect of surface pH of silica
by T. Cserháti and M. Szögyi (Budapest, Hungary) 249

Effective sample preparation method for extracting biologically active compounds from different matrices by a microwave technique
by K. Ganzler, I. Szinai and A. Salgó (Budapest, Hungary) 257

Polyalkylenes used as stationary phases for preparative reversed-phase liquid chromatography
by E. Krause and D. Smettan (Berlin, G.D.R.) and F. Loth and H. Herma (Teltow, G.D.R.) 263

Non-equilibrium analytical focusing field-flow fractionation using intrinsic hydrodynamic force and integral Doppler anemometry
by V. L. Kononenko and J. K. Shimkus (Moscow, U.S.S.R.) 271

Spacer-displacement and carrier spacer-displacement thin-layer chromatography
by H. Kalász (Budapest, Hungary) and M. Báthori and B. Matkovics (Szeged, Hungary) . 287

Preparative high-performance liquid chromatographic columns with an intra-column injection system
by B. Porsch, J. Voslář, J. Rosol and V. Kubánek (Prague, Czechoslovakia) 295

Study of the characteristics and analytical application of a capillary chromatograph with an electrochemical detector
by V. F. Ruban, I. A. Anisimova and B. G. Belen'kii (Leningrad, U.S.S.R.) 307

Peculiarities of aromatic hydrocarbon retention in normal-phase high-performance liquid chromatography with eluents containing halide derivatives
by S. N. Lanin and Yu. S. Nikitin (Moscow, U.S.S.R.) 315

Procedure for the purification of streptavidin by hydrophobic interaction chromatography
by W.-D. Schwidop, P. Klossek, R. Müller and R. Claus (Leipzig, G.D.R.) 325

Applications

High-performance liquid chromatographic method for the study of solvent effects on the peptidase and esterase activity of thermitase
by S. Dudek, S. Friebe and P. Hermann (Halle, G.D.R.) 333

Determination of cannabinoid acids by high-performance liquid chromatography of their neutral derivatives formed by thermal decarboxylation. I. Study of the decarboxylation process in open reactors
by T. Veress, J. I. Szanto and L. Leisztner (Budapest, Hungary) 339

Ion-exchange separation of inorganic anions on a HEMA 1000 Q-L column
by V. Pacáková, K. Štulík and W. Mingjia (Prague, Czechoslovakia) 349

Enantiomeric separation of racemic α-substituted α-methylbutyric acid derivatives on microcrystalline cellulose triacetate
by A. Kunath and R. Ozegowski (Berlin, G.D.R.) 361

Application of sodium lauryl sulphate ion-pairing reversed-phase high-performance liquid chromatography in the separation of some isomeric and other closely related basic compounds in pharmaceutical analysis
by M. Bidló-Oglóy (Budapest, Hungary) 367

High-performance liquid chromatographic method for direct resolution of the indobufen enantiomeric components
by G. Perrone and M. Farina (Milan, Italy) 373

Determination of N-(3-nitro-4-quinoline)morpholino-4-carboxamidine in plasma by high-performance liquid chromatography
by S. Benkő and J. Lengyel (Budapest, Hungary) 379

Aromatic ring-type analysis of complex hydrocarbon mixtures by high-performance liquid chromatography with photodiode-array detection and curve resolution
by H. Bünger (Berlin, G.D.R.), A. Meister (Gatersleben, G.D.R.) and M. Otto (Freiberg, G.D.R.) 383

Separation of hydroxylated polyenoic fatty acid enantiomers on Pirkle-type chiral phase high-performance liquid chromatographic columns
by H. Kühn and R. Wiesner (Berlin, G.D.R.) 391

Retention of some organic electrolytes in ion-pair reversed-phase high-performance liquid and reversed-phase high-performance thin-layer chromatographic systems
by M. L. Bieganowska, A. Petruczynik and M. Gadzikowska (Lublin, Poland) 403

Aldehyde and nucleotide separations by high-performance liquid chromatography. Application to phenylhydrazine-induced damage of erythrocytes and reticulocytes
by T. Grune, W. Siems, A. Werner and G. Gerber (Berlin, G.D.R.), M. Kostić (Kragujevac, Yugoslavia) and H. Esterbauer (Graz, Austria) 411

Ion-pair high-performance liquid chromatography of purine compounds in the small intestinal mucosa of children with coeliac disease
by Yu. V. Tikhonov, A. M. Pimenov, S. A. Uzhevko and R. T. Toguzov (Moscow, U.S.S.R.) 419

Use of chromatography for the preparation of homologous tetanus antitoxin
by V. A. Alyoshkin, V. N. Borisova, A. M. Ponomaryova and M. Y. Popkov (Moscow, U.S.S.R.) 425

Author Index 433